

**Supplementary Table 4.** Comparison of Adult and Pediatric Cohort in The Inflammatory Bowel Disease Registry

Variable	PIBD cohort CD	Adult cohort CD <sup>22</sup>	PIBD cohort UC	Adult cohort UC <sup>22</sup>
Male sex	62.8	61.2	63.7	61.4
Age at diagnosis (yr)	12.80±3.89	34.00±12.80	13.70±3.91	38.20±12.70
Diagnostic delay (mo)	23.56±27.76	42.2±46.1	19.37±26.76	37.20±43.80
Smoking	1.7	7.16	0.8	7.63
History of ATT	17.7	26.09	1.7	1.21
Family history	8.5	4.13	9.7	4.34
Disease location in CD				
L1 (ileal)	31.4	27.3		
L2 (colonic)	37.1	24.3		
L3 (ileocolonic)	26.3	40.9		
Isolated L4 (upper GI involvement)	5.1	5.0		
B1 (inflammatory)	57.1	67.8		
B2 (stricturing)	31.4	22.9		
B3 (fistulizing)	10.2	9.4		
Perianal disease	7.4	2.5		
Disease location in UC				
Proctitis			12.4	18.7
Proctosigmoiditis			15.0	51.0
Beyond splenic flexure			72.6	30.3
Treatment given (UC+CD)				
Steroids	69.7	39.7	56.6	39.5
Immunomodulators	59.4	45.9	38.9	34.6
Biologics	5.7	5.7	6.2	1.8

Value are presented as percent or mean ± standard deviation.

IBD, inflammatory bowel disease; PIBD, pediatric IBD; CD, Crohn's disease; UC, ulcerative colitis; ATT, anti-tubercular therapy; GI, gastrointestinal.